

Computational Statistics

A multinomial generalized linear mixed model for clustered competing risks data --Manuscript Draft--

Manuscript Number:	COST-D-22-00357R2
Full Title:	A multinomial generalized linear mixed model for clustered competing risks data
Article Type:	Original Paper
Keywords:	Cause-specific cumulative incidence function; Within-cluster dependence; Template Model Builder; Laplace approximation; Automatic differentiation
Manuscript Classifications:	1.01590: Generalized Linear Mixed Models; 1.02290: Likelihood Inference; 1.02570: Maximum Likelihood; 1.04140: Simulation Study; 1.04730: Survival Analysis
Corresponding Author:	Henrique Laureano Instituto de Pesquisa Pelé Pequeno Príncipe Curitiba, Paraná BRAZIL
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Instituto de Pesquisa Pelé Pequeno Príncipe
Corresponding Author's Secondary Institution:	
First Author:	Henrique Aparecido Laureano
First Author Secondary Information:	
Order of Authors:	Henrique Aparecido Laureano Ricardo Rasmussen Petterle Guilherme Parreira da Silva Paulo Justiniano Ribeiro Junior Wagner Hugo Bonat
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	Clustered competing risks data are a complex failure time data scheme. Its main characteristics are the cluster structure, which implies a latent within-cluster dependence between its elements, and its multiple variables competing to be the one responsible for the occurrence of an event, the failure. To handle this kind of data, we propose a full likelihood approach, based on generalized linear mixed models instead the usual complex frailty model. We model the competing causes in the probability scale, in terms of the cumulative incidence function (CIF). A multinomial distribution is assumed for the competing causes and censorship, conditioned on the latent effects that are accommodated by a multivariate Gaussian distribution. The CIF is specified as the product of an instantaneous risk level function with a failure time trajectory level function. The estimation procedure is performed through the R package TMB (Template Model Builder), an C++ based framework with efficient Laplace approximation and automatic differentiation routines. A large simulation study was performed, based on different latent structure formulations. The model fitting was challenging and our results indicated that a latent structure where both risk and failure time trajectory levels are correlated is required to reach reasonable estimation.

List of responses to the comments for the author of: A multinomial generalized linear mixed model for clustered competing risks data

March 26, 2023

Co-Editor

Based on the advice received, I have decided that your manuscript can be accepted for publication after you have carried out the corrections as suggested by the reviewer(s).

Author's response

We thanks the positive evaluation and we addressed in the paper the corrections and suggestions of the reviewers.

Reviewer 1

The authors have positively answered to all the issues arisen.

Reviewer 2

1. Please incorporate the comparison with He et al. (2022) discussed in the author's response into the paper (Introduction/Discussion).

Author's response

We thanks for the literature recommendation. The robust approach proposed by [He et al. \(2022\)](#) has been incorporated into the paper.

Reviewer 2

2. The authors mentioned that the Laplace-approximated MLE converges faster than the EM (which has a linear convergence rate), do we know at what rate it converges, e.g., approximately quadratic?

Author's response

We thanks fot the insightful comment. The Laplace approximation for the latent effects of a mixed model consists of two optimizations, an inner and an outer optimization. The inner one is made through a Newton-Raphson algorithm, Newton's method with a quadratic convergence rate. The external optimization is made through a Quasi-Newton Method, the BFGS for instance, which in our class of models has a superlinear convergence rate.

References

He, Y., Kim, S., Mao, L. and Ahn, K. W. (2022). Marginal semiparametric transformation models for clustered multivariate competing risks data, *Statistics in Medicine* **41**: 5349–5364.

A multinomial generalized linear mixed model for clustered competing risks data

Henrique Aparecido Laureano^{* 1}, Ricardo Rasmussen Petterle²,
Guilherme Parreira da Silva³, Paulo Justiniano Ribeiro Junior⁴, and
Wagner Hugo Bonat⁵

¹ Instituto de Pesquisa Pelé Pequeno Príncipe, Curitiba, Brasil

² Departamento de Medicina Integrada, Universidade Federal do Paraná,
Curitiba, Brasil

^{3,4,5} Laboratório de Estatística e Geoinformação, Departamento de
Estatística, Universidade Federal do Paraná, Curitiba, Brasil

March 26, 2023

Abstract

Clustered competing risks data are a complex failure time data scheme. Its main characteristics are the cluster structure, which implies a latent within-cluster dependence between its elements, and its multiple variables competing to be the one responsible for the occurrence of an event, the failure. To handle this kind of data, we propose a full likelihood approach, based on generalized linear mixed models instead the usual complex frailty model. We model the competing causes in the probability scale, in terms of the cumulative incidence function (CIF). A multinomial distribution is assumed for the competing causes and censorship, conditioned on the latent effects that are accommodated by a multivariate Gaussian distribution. The CIF is specified as the product of an instantaneous risk level function with a failure time trajectory level function. The estimation procedure is performed through the R package TMB (Template Model Builder), an C++ based framework with efficient Laplace approximation and automatic differentiation routines. A large simulation study was performed, based on different latent structure formulations. The model fitting was challenging and our results indicated that a latent structure where both risk and failure time trajectory levels are correlated is required to reach reasonable estimation.

^{*}E-mail: henriqueaparecidolaureano@gmail.com

Keywords: Cause-specific cumulative incidence function; Within-cluster dependence;
Template Model Builder; Laplace approximation; Automatic differentiation.

1 Introduction

Competing risks data, and more generally failure time data, can be modeled in two possible scales: the hazard and the probability scale, with the former being the most popular. A competing risks process can be seen as the multivariate extension of a failure time process, having multiple causes competing to be the one responsible for the desired event occurrence, properly, a failure. In Figure 1 a visual aid is provided considering m competing causes, where zero represents the initial state.




Figure 1: Illustration of competing risks process.

Failure time data is the branch of Statistics responsible to handle random variables describing the time until the occurrence of an event, a failure (Kalbfleisch and Prentice; 2002; Hougaard; 2000). The time until a failure is called survival experience, and is the modeling object. To accommodate the number of possible causes for a failure there is the competing risks data scheme. More specifically, its clustered version with groups of elements sharing some non-observed latent dependence structure.

When this framework is applied in real-world situations, we have to be able to handle with the nonoccurrence of the desired event, by any of the competing causes, for, let us say, *logistic reasons* (short-time study and outside scope causes are some examples). This, generally noninformative, nonoccurrence of the event is called censorship.

When the elements under study are organized in clusters (a family, e.g.), it opens space to what is called *family studies*. In family studies, the goal is to accommodate the non-observed latent dependence and try to understand the relationship between the family elements. In other words, how the occurrence of an event in a subject affects the survival experience for the same or similar event.

The survival experiences is usually modeled in the hazard (failure rate) scale, and with the latent within-cluster dependence accommodation we have what is called a frailty model (Clayton; 1978; Vaupel et al.; 1979; Liang et al.; 1995; Petersen; 1998). The use of frailty models implies in complicated likelihood functions and inference routines done via elaborated and slow EM algorithms (Nielsen et al.; 1992; Klein; 1992) or inefficient MCMC schemes (Hougaard; 2000). With multiple survival experiences, the general idea is the same but with even more elaborated likelihoods (Prentice et al.; 1978; Therneau and Grambsch; 2000) or mixture model approaches (Larson and Dinse; 1985; Kuk; 1992).

When in the hazard scale, the interpretations are in terms of hazard rates. A less usual scale but with a more appealing interpretation is the probability scale. For competing risks data, the work on the probability scale is done by means of the cumulative incidence function (CIF) (Andersen et al.; 2012), with the main modeling approach being the subdistribution (Fine and Gray; 1999). In family studies or populational studies, where the main interest relies on describing and understanding, in terms of regressors, the long-run behavior of a process, the CIF is a suitable modeling choice having the bonus of the interpretation in terms of the probability scale.

For clustered competing risks data there are some available options but with a lack of predominance. The options vary in terms of likelihood specification, with its majority being designed for bivariate CIFs, where increasing the CIF's dimension is a limitation. Some of the existing options are (i) nonparametric approaches (Cheng et al.; 2007, 2009); (ii) linear transformation models (Fine; 1999; Gerds et al.; 2012; He et al.; 2022); (iii) semiparametric approaches based on composite likelihoods (Shih and Albert; 2009; Cederkvist et al.; 2019), estimating equations (Scheike and Sun; 2012; Cheng and Fine; 2012), copulas (Scheike et al.; 2010), or mixtures (Naskar et al.; 2005; Shi et al.; 2013).

Besides the interpretation, by modeling the CIF it is possible to specify complex within-cluster dependence structures. We follow Cederkvist et al. (2019) and work with a CIF specification based on its decomposition in instantaneous risk and failure time trajectory functions, with both being cluster-specifics and possible correlated. Instead of an elaborated composite likelihood approach, as a modeling framework, we use a generalized linear mixed model (GLMM) specification. Through a GLMM we have a straightforward full likelihood specification, easy to virtually extend to any number of competing causes, and capable to allow for complex CIF structures. To make the estimation and inferential process the most efficient as possible we take advantage of state-of-art computational libraries and efficiently implemented routines under the TMB (Kristensen et al.; 2016) package of the R (R Core Team; 2021) statistical software.

The class of generalized linear models (GLMs) (Nelder and Wedderburn; 1972) is probably the most popular statistical modelling framework. Despite its flexibility, the GLMs are not suitable for dependent data. For the analysis of such data, Laird and Ware (1982) proposed the random effects regression models for longitudinal/repeated-measures data, and Breslow and Clayton (1993) presented the GLMMs for the analysis of non-Gaussian outcomes. In this framework, we can accommodate all competing causes of failure and censorship under a multinomial probability distribution. The latent within-cluster dependence is accommodated via a multivariate normal distribution, and the cause-specific CIFs via the model's link function.

The main goal of this article is to propose a likelihood approach for parameter estimation and inference in the context of clustered competing risks data with a flexible within-cluster dependence structure. The model specification is based on the approach of

Cederkvist et al. (2019) which in turn resembles the GLMM framework. However, for parameter estimation, we propose a full likelihood approach where the marginal likelihood function is obtained using the Laplace approximation. On the other hand, Cederkvist et al. (2019) employed an intricate composite likelihood combined with Adaptive Gaussian Quadrature. The main advantage of our approach is the computational efficiency allowing us to use state-of-art numerical libraries for automatic differentiation and parallel computation.

The main contributions of this article are: (i) introducing the modeling of cause/cluster-specific CIFs of clustered competing risks data into an efficient implementation of the GLMMs framework; (ii) performing a extensive simulation study to check the properties of the maximum likelihood estimator to learn the cause-specific CIF forms and the feasibility of the within-cluster dependence structure.; (iii) providing R code and C++ implementation for the used GLMMs.

The work is organized as follows. Section 2 presents the CIF specification and the multinomial GLMM. Section 3 presents the estimation and inferential routines. Section 4 presents the performed simulation studies to check the model viability. Finally, the main contributions of the article are discussed in Section 5.

2 Model

2.1 Cluster-specific cumulative incidence function (CIF)

Consider that the observed follow-up time of a subject is given by $T = \min(T^*, C)$, where T^* denote the failure time and C denote the censoring time. Given the possible covariates $\mathbf{x}$, for a cause-specific of failure k , the cumulative incidence function (CIF) is defined as

$$\begin{aligned} F_k(t | \mathbf{x}) &= \mathbb{P}[T \leq t, K = k | \mathbf{x}] = \int_0^t f_k(z | \mathbf{x}) \, dz \\ &= \int_0^t \lambda_k(z | \mathbf{x}) S(z | \mathbf{x}) \, dz, \quad t > 0, \quad k = 1, \dots, K, \end{aligned}$$

where $f_k(t | \mathbf{x})$ is the (sub)density for the time to a type k failure. The subdensity is composed by the cause-specific hazard function or process $\lambda_k(t | \mathbf{x})$, representing the instantaneous rate for failures of type k at time t given $\mathbf{x}$ and all other failure types (competing causes). If we sum up all cause-specific hazard functions we get the overall hazard function $\lambda(t | \mathbf{x})$. From the overall hazard function we arrive in the overall survival function,

$$S(t | \mathbf{x}) = \mathbb{P}[T > t | \mathbf{x}] = \exp \left\{ - \int_0^t \lambda(z | \mathbf{x}) \, dz \right\},$$

the second function that compose the subdensity $f_k(t | \mathbf{x})$. A comprehensive reference for all these definitions is the book of [Kalbfleisch and Prentice \(2002\)](#).

To take into consideration our clustered/family structure, we follow the same CIF specification of [Cederkvist et al. \(2019\)](#). For two competing causes of failure, the cause-specific CIFs are specified in the following manner

$$F_k(t | \mathbf{x}, u_1, u_2, \eta_k) = \underbrace{\pi_k(\mathbf{x}, u_1, u_2)}_{\text{cluster-specific risk level}} \times \underbrace{\Phi[w_k g(t) - \mathbf{x}\boldsymbol{\gamma}_k - \eta_k]}_{\text{cluster-specific failure time trajectory}}, \quad t > 0, \quad k = 1, 2, \quad (1)$$

i.e., as the product of a cluster-specific risk level with a cluster-specific failure time trajectory, resulting in a cluster-specific CIF. What makes these components cluster-specific are $\mathbf{u} = \{u_1, u_2\}$ and $\boldsymbol{\eta} = \{\eta_1, \eta_2\}$, Gaussian distributed latent effects with zero mean and potentially correlated,

$$\begin{bmatrix} u_1 \\ u_2 \\ \eta_1 \\ \eta_2 \end{bmatrix} \sim \text{Multivariate Normal} \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{u_1}^2 & \text{cov}(u_1, u_2) & \text{cov}(u_1, \eta_1) & \text{cov}(u_1, \eta_2) \\ & \sigma_{u_2}^2 & \text{cov}(u_2, \eta_1) & \text{cov}(u_2, \eta_2) \\ & & \sigma_{\eta_1}^2 & \text{cov}(\eta_1, \eta_2) \\ & & & \sigma_{\eta_2}^2 \end{bmatrix} \right).$$

The cluster-specific risk levels are modeled by a multinomial logistic regression model with latent effects, i.e.

$$\pi_k(\mathbf{x}, \mathbf{u}) = \frac{\exp\{\mathbf{x}\boldsymbol{\beta}_k + u_k\}}{1 + \exp\{\mathbf{x}\boldsymbol{\beta}_1 + u_1\} + \exp\{\mathbf{x}\boldsymbol{\beta}_2 + u_2\}}, \quad k = 1, 2, \quad (2)$$

where the $\boldsymbol{\beta}_k$'s are the coefficients responsible for quantifying the impact of the covariates in the cause-specific risk levels. For individuals from the same cluster/family, at the same time point, the $\boldsymbol{\beta}_k$ s have the well-known odds ratio interpretation.

The second component of [Equation 1](#) is the cluster-specific failure time trajectory

$$\Phi[w_k g(t) - \mathbf{x}\boldsymbol{\gamma}_k - \eta_k], \quad t > 0, \quad k = 1, 2,$$

where $\Phi(\cdot)$ is the cumulative distribution function of a standard Gaussian distribution. With regard to the function $g(t)$, it plays a crucial role since the proposed CIF separation is only possible with it. It is used a time t transformation given by

$$g(t) = \text{arctanh} \left(\frac{t - \delta/2}{\delta/2} \right), \quad t \in (0, \delta), \quad g(t) \in (-\infty, \infty),$$

where δ depends on the data and cannot exceed the maximum observed follow-up time τ , i.e. $\delta \leq \tau$. With this Fisher-based transformation the value of the cluster-specific failure time trajectory is equal 1 at time δ . Consequently, $F_k(\delta | \mathbf{x}, \mathbf{u}, \eta_k) = \pi_k(\mathbf{x} | \mathbf{u})$ and

we can interpret $\pi_1(\mathbf{x} | \mathbf{u})$ and $\pi_2(\mathbf{x} | \mathbf{u})$ as the cause-specific cluster-specific risk levels at time δ . The cluster-specific survival function is given by $S(t | \mathbf{x}, \mathbf{u}, \boldsymbol{\eta}) = 1 - F_1(t | \mathbf{x}, \mathbf{u}, \eta_1) - F_2(t | \mathbf{x}, \mathbf{u}, \eta_2)$.

A direct understanding of all coefficients/parameters in Equation 1 can be reached via the top-placed illustrations in Figure 2. We consider two competing causes, without covariates, and plot the cluster-specific CIF of just one failure cause. We see that:

- The β 's are related to the curve's maximum value, i.e. bigger the β highest the CIF;
- The γ_k 's are the coefficients responsible for quantifying the impact of the covariates in the cause-specific failure time trajectories, i.e. the shape of the cumulative incidence. We see that the γ 's are also related with an idea of midpoint and consequently, growth speed. The fact that γ_k enters negatively in the cluster-specific failure time trajectory makes that a negative value causes an advance towards the curve, whereas a positive value causes a delay;
- Last but not least, the w 's. With negative values we have a decreasing curve and with positive values an increasing curve, the behavior of interest.

Cluster-specific Cumulative Incidence Function (CIF)




Figure 2: Curve behaviors for different parameter settings, showing then the corresponding parameter effects in a cluster-specific cumulative incidence function (CIF).

Remains to talk about the within-cluster dependence induced by the latent effects in $\mathbf{u}$ and $\boldsymbol{\eta}$. To help in the discussion, the bottom-plots on [Figure 2](#) illustrates the cluster-specific CIF for a given failure cause in a model without covariates, let us call it failure cause 1 (in total we have two). The latent effects u_1 and u_2 always appear together in the cluster-specific risk level, as consequence they have a joint effect on the cumulative incidence of both causes. As we can see in [Figure 2](#), an increase in u_k will increase the risk of failure from cause k . The interpretation of $\text{cov}(\eta_1, \eta_2)$ and $\text{cov}(u_1, u_2)$ is straightforward, and those values are in most of the cases positive, as said in [Cederkvist et al. \(2019\)](#). With regard to $\text{cov}(u_k, \eta_k)$, negative values are the common situation. A negative correlation between η_k and u_k imply that when η_k decreases, u_k increases and conversely when η_K increases, u_k decreases. In other words, an increased risk level is reached quickly and a decreased risk level is reached later, respectively. With regard to cross-cause correlation between η and u , positive values are the common situation where late onset of one failure cause is associated with a high absolute risk of another failure cause.

2.2 Model specification

The generalized linear mixed model (GLMM) for clustered competing risks data is specified in the following hierarchical fashion. By simplicity, we focus on two competing causes of failure but an extension is straightforward. For two competing causes of failure, the failure or censorship, y , of a subject i , in the cluster/family j and time t , is modeled by

$$y_{ijt} \mid \{u_{1j}, u_{2j}, \eta_{1j}, \eta_{2j}\} \sim \text{Multinomial}(p_{1ijt}, p_{2ijt}, p_{3ijt})$$

$$\begin{bmatrix} u_1 \\ u_2 \\ \eta_1 \\ \eta_2 \end{bmatrix} \sim \text{MN} \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{u_1}^2 & \text{cov}(u_1, u_2) & \text{cov}(u_1, \eta_1) & \text{cov}(u_1, \eta_2) \\ & \sigma_{u_2}^2 & \text{cov}(u_2, \eta_1) & \text{cov}(u_2, \eta_2) \\ & & \sigma_{\eta_1}^2 & \text{cov}(\eta_1, \eta_2) \\ & & & \sigma_{\eta_2}^2 \end{bmatrix} \right)$$

$$\begin{aligned} p_{kijt} &= \frac{\partial}{\partial t} F_k(t \mid \mathbf{x}, u_1, u_2, \eta_k) \\ &= \frac{\exp\{\mathbf{x}_{kij}\boldsymbol{\beta}_k + u_{kj}\}}{1 + \sum_{m=1}^{K-1} \exp\{\mathbf{x}_{mij}\boldsymbol{\beta}_m + u_{mj}\}} \\ &\quad \times w_k \frac{\delta}{2\delta t - 2t^2} \phi \left(w_k \text{arctanh} \left(\frac{t - \delta/2}{\delta/2} \right) - \mathbf{x}_{kij}\boldsymbol{\gamma}_k - \eta_{kj} \right), \\ k &= 1, 2. \end{aligned} \tag{3}$$

The probabilities are given by the derivative w.r.t. time t of the cluster-specific CIF. The choice of a multinomial logistic regression model ensures that the sum of all predicted

cause-specific CIFs does not exceed 1. Considering two competing causes of failure, we have a multinomial with three classes. The third class exists to handle the censorship and its probability is given by the complementary to reach 1. To better describe these curve behaviors we have [Figure 3](#).




Figure 3: Cluster-specific cumulative incidence function (CIF) curves and their derivatives (dCIF) for a random scenario with two competing causes.

This framework in [Equation 3](#) results in what we call multiGLMM, a multinomial GLMM to handle the CIF of clustered competing risks data. For a random sample, the corresponding marginal likelihood function is given by

$$\begin{aligned}
L(\boldsymbol{\theta} ; y) &= \prod_{j=1}^J \int_{\mathbb{R}^4} \pi(y_j | \mathbf{r}_j) \times \pi(\mathbf{r}_j) d\mathbf{r}_j \\
&= \prod_{j=1}^J \int_{\mathbb{R}^4} \underbrace{\left\{ \prod_{i=1}^{n_j} \prod_{t=1}^{n_{ij}} \left(\frac{(\sum_{k=1}^K y_{kijt})!}{y_{1ijt}! y_{2ijt}! y_{3ijt}!} \prod_{k=1}^K p_{kijt}^{y_{kijt}} \right) \right\}}_{\text{fixed effect component}} \times \\
&\quad \underbrace{(2\pi)^{-2} |\Sigma|^{-1/2} \exp \left\{ -\frac{1}{2} \mathbf{r}_j^\top \Sigma^{-1} \mathbf{r}_j \right\}}_{\text{latent effect component}} d\mathbf{r}_j \\
&= \prod_{j=1}^J \int_{\mathbb{R}^4} \underbrace{\left\{ \prod_{i=1}^{n_j} \prod_{t=1}^{n_{ij}} \prod_{k=1}^K p_{kijt}^{y_{kijt}} \right\}}_{\text{fixed effect}} \underbrace{(2\pi)^{-2} |\Sigma|^{-1/2} \exp \left\{ -\frac{1}{2} \mathbf{r}_j^\top \Sigma^{-1} \mathbf{r}_j \right\}}_{\text{latent effect component}} d\mathbf{r}_j, \quad (4)
\end{aligned}$$

where $\boldsymbol{\theta} = [\boldsymbol{\beta} \ \boldsymbol{\gamma} \ \mathbf{w} \ \boldsymbol{\sigma}^2 \ \boldsymbol{\rho}]^\top$ is the parameters vector to be maximized. In our framework, a subject can fail from just one competing cause or get censor, at a given time. Thus, the fraction of factorials in the fixed effect component is made only by 0's and 1's. The matrix Σ is the variance-covariance matrix, which parameters are given by $\boldsymbol{\sigma}^2$ and $\boldsymbol{\rho}$.

To each cluster j we have a product of two components. The fixed effect component, given by a multinomial distribution with its probabilities specified through the cluster-

specific CIF (Equation 1) and, the latent effect component, given by a multivariate Gaussian distribution. To each subject i that composes a cluster j we have its specific fixed effects contribution. The likelihood in Equation 4 is the most general as possible, allowing for repeated measures to each subject. Since all subjects of a given cluster shares the same latent effect, we have just one latent effect contribution multiplying the product of fixed effect contributions. As we do not observe the latent effect variables, $\mathbf{r}_j$, we integrate out in it. With two competing causes of failure we have four latent effects, a multivariate Gaussian distribution in four dimensions. Consequently, for each cluster, we approximate an integral in four dimensions. The product of these approximated integrals results in the called marginal likelihood, to be maximized in $\boldsymbol{\theta}$.

3 Estimation and inference

Our goal is to estimate the parameter vector $\boldsymbol{\theta} = [\boldsymbol{\beta} \ \boldsymbol{\gamma} \ \mathbf{w} \ \boldsymbol{\sigma}^2 \ \boldsymbol{\rho}]^\top$. The likelihood for $\boldsymbol{\theta}$ can be written as

$$L(\boldsymbol{\theta} \mid \mathbf{y}, \mathbf{u}) = \prod_{i=1}^I \prod_{j=1}^{n_i} f(y_{ij} \mid \mathbf{u}_i, \boldsymbol{\beta}, \boldsymbol{\Sigma}) f(\mathbf{u}_i \mid \boldsymbol{\Sigma}). \quad (5)$$

From standard probability theory is easy to see that in the right-hand side (r.h.s.) we have a joint density, consequently, Equation 5 represents what is called a full or joint likelihood function. The latent effect $\mathbf{u}$ is *latent*, i.e. we do not observe it. To handle this we use the Laplace approximation.

If we have a joint density we can just integrate out the undesired variable resulting in

$$\begin{aligned} L(\boldsymbol{\theta} \mid \mathbf{y}) &= \prod_{i=1}^I \int_{\mathcal{R}^{\mathbf{u}_i}} \left[\prod_{j=1}^{n_i} f(y_{ij} \mid \mathbf{u}_i, \boldsymbol{\beta}, \boldsymbol{\Sigma}) f(\mathbf{u}_i \mid \boldsymbol{\Sigma}) \right] d\mathbf{u}_i \\ &= \prod_{i=1}^I \int_{\mathcal{R}^{\mathbf{u}_i}} f(\mathbf{y}_i, \mathbf{u}_i \mid \boldsymbol{\theta}) d\mathbf{u}_i, \end{aligned} \quad (6)$$

a marginal density that keeps the parameters $\{\boldsymbol{\sigma}^2 \ \boldsymbol{\rho}\}$ of the integrated variable. To handle this integration step a clever choice is to take advantage of the exponential family structure together with the Gaussian latent effects distribution. These ideas converge to an adaptive Gaussian quadrature with one integration point, also known as *Laplace approximation* (Molenberghs and Verbeke; 2005; Shun and McCullagh; 1995; Tierney and Kadane; 1986; Wood; 2015).

We may approximate an analytically intractable integral in a way to obtain a tractable closed-form expression allowing the numerical maximization of the resulting marginal likelihood function (Bonat and Ribeiro Jr; 2016). The Laplace approximation has been

designed to approximate integrals in the form

$$\int_{\mathcal{R}^{\mathbf{u}_i}} \exp\{Q(\mathbf{u}_i)\} d\mathbf{u}_i \approx (2\pi)^{n_{\mathbf{u}}/2} |Q''(\hat{\mathbf{u}}_i)|^{-1/2} \exp\{Q(\hat{\mathbf{u}}_i)\}, \quad (7)$$

where $Q(\mathbf{u}_i)$ is a known unimodal bounded function, and $\hat{\mathbf{u}}_i$ is the value for which $Q(\mathbf{u}_i)$ is maximized. A advantage of the Laplace approximation approach in a GLMM is the exponential family structure. In a usual GLMM the response follows a one-parameter exponential family distribution that can be written as

$$f(\mathbf{y}_i | \mathbf{u}_i, \boldsymbol{\theta}) = \exp \left\{ \mathbf{y}_i^\top (\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i) - \mathbf{1}_i^\top b(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i) + \mathbf{1}_i^\top c(\mathbf{y}_i) \right\},$$

where $b(\cdot)$ and $c(\cdot)$ are known functions. This general and easy to compute expression together with a (multivariate) Gaussian distribution, highlights the convenience of the Laplace method. The $Q(\mathbf{u}_i)$ function to be maximized can be expressed as

$$\begin{aligned} Q(\mathbf{u}_i) &= \mathbf{y}_i^\top (\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i) - \mathbf{1}_i^\top b(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i) + \mathbf{1}_i^\top c(\mathbf{y}_i) \\ &\quad - \frac{n_{\mathbf{u}}}{2} \log(2\pi) - \frac{1}{2} \log |\boldsymbol{\Sigma}| - \frac{1}{2} \mathbf{u}_i^\top \boldsymbol{\Sigma}^{-1} \mathbf{u}_i. \end{aligned} \quad (8)$$

The approximation in Equation 7 requires the maximum $\hat{\mathbf{u}}_i$ of the function $Q(\mathbf{u}_i)$. As we assume a Gaussian distribution with a known mean for the latent effects, we have the perfect initial guess for a Hessian-based maximization method, as the Newton-Raphson (NR) algorithm. Bonat and Ribeiro Jr (2016) presents the generic expressions for the derivatives required by the NR method, given by the following:

$$\begin{aligned} Q'(\mathbf{u}_i^{(k)}) &= \{\mathbf{y}_i - b'(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i^{(k)})\}^\top - \mathbf{u}_i^{(k)\top} \boldsymbol{\Sigma}^{-1}, \\ Q''(\mathbf{u}_i^{(k)}) &= -\text{diag}\{b''(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \mathbf{u}_i^{(k)})\} - \boldsymbol{\Sigma}^{-1}. \end{aligned}$$

The marginal log-likelihood function returned by the Laplace approximation, to each individual or unit under study i , is as follows:

$$\begin{aligned} l(\boldsymbol{\theta} | \mathbf{y}_i) &= \log L(\boldsymbol{\theta} | \mathbf{y}_i) = \frac{n}{2} \log(2\pi) - \frac{1}{2} \log |\text{diag}\{b''(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \hat{\mathbf{u}}_i)\} + \boldsymbol{\Sigma}^{-1}| \\ &\quad + \mathbf{y}_i^\top (\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \hat{\mathbf{u}}_i) - \mathbf{1}_i^\top b(\mathbf{x}_i \boldsymbol{\beta} + \mathbf{z}_i \hat{\mathbf{u}}_i) + \mathbf{1}_i^\top c(\mathbf{y}_i) \\ &\quad - \frac{n_{\mathbf{u}}}{2} \log(2\pi) - \frac{1}{2} \log |\boldsymbol{\Sigma}| - \frac{1}{2} \hat{\mathbf{u}}_i^\top \boldsymbol{\Sigma}^{-1} \hat{\mathbf{u}}_i, \end{aligned}$$

that can now be numerically maximized over the model parameters $\boldsymbol{\theta} = [\boldsymbol{\beta} \ \boldsymbol{\gamma} \ \mathbf{w} \ \boldsymbol{\sigma}^2 \ \boldsymbol{\rho}]^\top$ using a quasi-Newton method as the Broyden-Fletcher-Goldfarb-Shanno (BFGS) algorithm (Nocedal and Wright; 2006) or the PORT routines (Dennis et al.; 1981; Gay; 1990), all available in the R (R Core Team; 2021) statistical software.

We use an efficient Laplace approximation routine implemented in TMB (Kristensen

et al.; 2016). Besides state-of-art linear algebra libraries and the possibility of performing the computations in parallel, TMB also offers an efficient implementation of automatic differentiation (Nocedal and Wright; 2006; Wood; 2015; Peyré; 2020), the state-of-art in gradients computation. In TMB the user should write its likelihood function in a C++ template file and then load it on R. This last characteristic makes TMB general (a vast range of models being allowed to be written) and powerful (low-level C++ model implementation).

4 Simulation studies

To verify the practical viability of our model we performed an extensive simulation study. As main complicator factors, we may mention the competing causes imbalance, where we typically have much more censorships than actual failures; and the high dimensionality problem, having a considerable number of parameters to estimate in both fixed and latent effects layers.

One of the main questions surrounding our model that we tried to tack down in this study was, if even when in the possession of a high correlated sample in the latent field, we are able to estimate all variance and covariance parameters. To answer this question we worked with four different models, with each one differentiating from the other in terms of latent effects structure. We had a model with (i) latent effects only on the risk level; (ii) only on the failure time trajectory level; (iii) on both levels but without cross-correlations, called a block-diag model; and (iv) a model with all possible correlations presented, called as a complete model. In Figure 4 a visual representation of these latent effects structures is presented, in terms of the corresponding variance-covariance matrices.




Figure 4: Simulation study variance-covariance matrix model designs, considering two competing causes of failure and consequently, four latent effects.

Besides the four latent effects structures, we worked with two CIF configurations

High CIF configuration : $\{\beta_1 = -2, \beta_2 = -1.5, \gamma_1 = 1, \gamma_2 = 1.5, w_1 = 3, w_2 = 4\}$;

Low CIF configuration : $\{\beta_1 = 3, \beta_2 = 2.5, \gamma_1 = 2.6, \gamma_2 = 4, w_1 = 5, w_2 = 10\}$.

Those two sets of parameter values together with the following variance-covariance structure values arise in CIF curves with peaks around 0.15 (low incidence scenario) and 0.60 (high incidence scenario). For numerical efficiency reasons, we worked with the log-variances and the Fisher-z-transformation of the correlation parameters.

$$\begin{aligned} \sigma_{u_1}^2 &= 1 \\ \sigma_{u_2}^2 &= 0.7 \\ \sigma_{\eta_1}^2 &= 0.6 \\ \sigma_{\eta_2}^2 &= 0.9, \end{aligned} \quad \text{Correlation structure} = \begin{pmatrix} u_1 & u_2 & \eta_1 & \eta_2 \\ 1 & 0.1 & -0.5 & 0.3 \\ & 1 & 0.3 & -0.4 \\ & & 1 & 0.2 \\ & & & 1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \eta_1 \\ \eta_2 \end{pmatrix}.$$

We also considered three cluster sizes (2, 5, and 10) and three sample sizes (5000, 30000, and 60000). We had 72 scenarios (4 latent effects structures $\times$ 2 CIF configurations $\times$ 3 cluster sizes $\times$ 3 sample sizes). For each scenario, we simulated 500 samples. In total, 36000 (72×500) model fittings. In the smallest scenario, we had 500 clusters, in the biggest one, 30000 clusters. The number of clusters is the number of Laplace approximations to be performed. With two competing causes of failure, we have a model with 16 parameters (6 in the fixed effect layer, 3 for each competing cause; and 10 in the latent effects layer, 4 variances, and 6 covariance parameters).

All models were run, in a parallelized fashion, in one of the two following Linux systems:

System 1 12 Intel(R) Core(TM) i7-8750H CPU @2.20GHz processors with 16GB RAM;

System 2 30 Intel(R) Xeon(R) CPU E5-2690v2 @3.00GHz processors and 206GB RAM.

The non-complete models (involving 2D Laplace approximations) are kind of fast, taking always less than 5 minutes of processing. In the most expensive scenarios (30K 4D Laplaces), the complete model takes 30 minutes. In terms of parameters estimation, the non-complete models fail to learn the data. They appear to be not structured enough to capture the data characteristics, showing that a full correlated latent field is really the most appropriate choice.

In the supplementary materials, we have several graphs summarizing the parameters estimate bias. In each figure, we have the estimate bias and its uncertainty described by a Wald-based confidence interval, i.e. ± 1.96 the bias standard deviation. We chose to use this uncertainty representation based on the point estimates instead of the standard error computations, since in several scenarios the model fails to compute all the standard errors, caused by Hessian numerical instabilities.

When we assume a non-zero cross-correlation structure (complete model), the improvements in terms of bias reduction are huge when compared with the non-complete models. The mean biases get very close to zero, the standard deviations decrease 50% or more, compared with the non-complete models, and the high CIF scenarios are the ones

with a much smaller bias-variance. All this is accomplished through the consideration of the cross-correlations.


In the *simpler* models, with a latent structure just in one level, is hard to see some significant difference between the clusters and sample sizes. With the complete model, in the other hand, the difference is clear: as we increase the clusters and the sample sizes, the bias-variance decreases. The mean-bias is basically always the same. The biggest bias-variances are obtained in the log-variances. A final remark to be made is about convergences. With the simpler models, not all of them work, having in some scenarios (generally the ones with 60 thousand data points) a 50~60% convergence rate. With the complete model, basically, almost all fits reach convergence (~95% performance).

About the implied mean-CIF curves, in [Figure 5](#) we have the high CIF scenario curves and in [Figure 6](#) the low CIF scenario curves. Since for all models we have a latent structure for the within-cluster dependency, the inherent idea is that this also affect the fixed-effect parameter estimates. By taking its average in each of the seventy-two scenarios, we are able to construct the mean CIF curves.

HIGH CIF SCENARIO

CIF of failure cause 1

True curve in dashed black



CIF of failure cause 2

True curve in dashed black





Figure 5: High cumulative incidence function (CIF) scenario curves. **cs** means the cluster size (2, 5, and 10), and 5, 30, and 60K means the sample size.

In Figure 5 is clear that with the complete model we get a perfect fit in all scenarios. The risk and time models estimate well the curve shape parameters but they fail to learn the max incidence. A compensation between curves is clear. In the low CIF scenarios in Figure 6, the estimation is clearly more difficult. The overall fits were bad. For the failure cause 2, the estimation quality is not so bad. The problem is when we look to the failure cause 1. The best joint fit is still with the complete model.

LOW CIF SCENARIO

CIF of failure cause 1

True curve in dashed black



CIF of failure cause 2

True curve in dashed black




Figure 6: Low cumulative incidence function (CIF) scenario curves. **cs** means the cluster size (2, 5, and 10), and 5, 30, and 60K means the sample size.

5 Discussion

The general goal of this paper was the proposition and evaluation of a maximum likelihood estimation approach for the analysis of clustered competing risks data. Focused on the probability scale, by means of the cumulative incidence function (CIF), instead of the usual hazard scale (Kalbfleisch and Prentice; 2002). We modeled the clustered competing risks on a latent-effects framework, a generalized linear mixed model (GLMM) (McCulloch and Searle; 2001), with a multinomial distribution for the competing risks and censorship, conditioned on the latent-effects. The within-cluster latent dependency was accommodated by a multivariate Gaussian distribution.

1 The failures by the competing causes and their respective censorships are modeled by
2 means of the CIF (Kalbfleisch and Prentice; 2002; Andersen et al.; 2012). The CIF was
3 accommodated in our GLMM framework in terms of the link function (McCullagh and
4 Nelder; 1989), as the product of two functions, one responsible to model the instantaneous
5 risk and the other the failure time trajectory, both in a cluster-specific fashion. This
6 particular GLMM formulation is what makes our model, particular. Thus, we have what
7 we call a multiGLMM: a multinomial GLMM for clustered competing risks data.

8 The two-function product CIF formulation was taken from Cederkvist et al. (2019)
9 but there they use a different estimation framework, a composite likelihood framework
10 (Lindsay; 1988; Cox and Reid; 2004; Varin et al.; 2011). Here we do a full likelihood
11 analysis instead. A composite approach is generally used when a full likelihood approach
12 is impossible or computationally impracticable. Our goal here was to assess a full like-
13 lihood framework taking advantage of state-of-the-art computational libraries together
14 with efficient algorithm implementations. We have all this with the R (R Core Team;
15 2021) package TMB (Kristensen et al.; 2016).

16 The applications in focus here were family studies. Besides the within-cluster/family
17 dependence, this kind of study is characterized by involving big samples, generally, pop-
18 ulations. Also, generally having a high number of small clusters, families, and a lot
19 of censorship. A maximum likelihood approach with the use of efficiently implemented
20 Laplace approximations (Tierney and Kadane; 1986; Bonat and Ribeiro Jr; 2016) together
21 with an automatic differentiation (AD) (Wood; 2015; Nosedal and Wright; 2006) routine,
22 all via TMB, is able to efficiently handle with a high number of clusters, independent of its
23 size. The multinomial distribution assumption, on its own, is an excellent probabilistic
24 choice since it can accommodate virtually any number of competing causes of failure and
25 its censorship. The presence of those two characteristics in our multiGLMM makes it an
26 efficient and scalable modeling framework for clustered competing risks data.

27 Even with our modeling framework being virtually able to handle any number of
28 competing causes of failure, we restrained ourselves to work here with only two of them.
29 With two competing causes, we have a 4×4 covariance matrix for the latent effects,
30 which implies ten covariance-matrix parameters, which is already a lot of parameters
31 to be estimated in a latent structure. Since our goal was to assess the viability of the
32 maximum likelihood estimation method, we kept it with two causes.

33 The so-called complete model, with latent effects on both risk and failure time tra-
34 jectory levels with cross-correlations, presented the best results. In its biggest scenario,
35 with 60000 data points and clusters of size 2, i.e. 30000 four-dimension integral approxi-
36 mations, the model fitting took 30 minutes, in parallel. Before TMB, we did a complete R
37 implementation writing our own Laplace approximation (Bonat and Ribeiro Jr; 2016) and
38 Newton-Raphson gradients and Hessian, by hand. In a scenario with 20000 data points
39 and clusters of size 2, it took around 30 hours, in parallel. In summary, by using TMB we

were able to increase the model size 3 times and to decrease the computational time 60 times. An incredible performance gain.

Also with the complete model, we performed a Bayesian analysis via `tmbstan` (Monahan and Kristensen; 2018). `tmbstan` enables MCMC sampling (Gelfand and Smith; 1990; Diaconis; 2009) from a TMB model object using Stan (Stan Development Team; 2019, 2020). Sampling was performed based on the probably state-of-art MCMC sampler algorithm, a Hamiltonian Monte Carlo (HMC) algorithm with the No-U-Turn Sampler (NUTS) extension (Hoffman and Gelman; 2014). We performed just one Bayesian model fitting in a modest scenario with 5000 data points and clusters of size 2. It took around 1 whole week of parallelized processing. The results were basically the same as the ones obtained with TMB but this high computational time just reinforces the MCMC framework limitation.

We proposed an ideally complete latent-effects formulation. The main underlying idea of the simulation study was to see in which scenarios we would be able to learn all the involved mean and covariance parameters. As part of that, simpler formulations were proposed. As result, we got that latent effects only in the risk level did not work. The optimization appears to got lost as if something was missing. Inserting latent effects only in the failure time trajectory level returned better results, but still not satisfactorily good. In most of the evaluated scenarios, the block-diagonal model appeared to be in the middle of them, as a compromise. The best results (smallest parameter estimates biases) were obtained with the complete model. The complete model works fine, mainly in the scenarios of high CIF configuration, and also as expected, as the sample size increases.

As an alternative to our model, we have two possible paths. We could instead of a conditional modeling framework (GLMM/latent-effects model), employ a marginal modeling framework. In this framework, instead of caring about the specification of a probability distribution to the competing causes conditioned on the latent effects, we just care about the specification of a mean and a variance structure. This approach does not have a likelihood function per se, but the estimation procedure tends to be easier than with the GLMM one. Marginal modeling frameworks that can be used here are the semiparametric linear transformation model (He et al.; 2022), also with the advantage of the non-necessity of modeling the censoring distribution, and the multivariate covariance generalized linear model (McGLM) (Bonat and Jørgensen; 2016; Bonat; 2018). How to exactly model the CIF of clustered competing risks data in this framework, is something to still be figured out.

The other path is by the use of a different way of modeling the dependence structure. Instead of a latent-effects approach, we could use copulas (Embrechts; 2009; Scheike et al.; 2010; Masarotto and Varin; 2012; Krupskii and Joe; 2013). How to do that is something to still be figured out in terms of which kind (conditional or marginal) and version (Archimedean-, Gauss-, Maltesian-, t -, hyperbolic-, zebra-, and elliptical-) of copula to

use, besides the estimability issue.

The C++ and R code for fitting our multiGLMM is available at <http://leg.ufpr.br/~henrique/papers/multiglmm/>.

Supplementary material




Figure 7: Parameter β_1 bias with ± 1.96 standard deviations. **cs** means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: β_2

with ± 1.96 standard deviations




Figure 8: Parameter β_2 bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: γ_1

with ± 1.96 standard deviations




Figure 9: Parameter γ_1 bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings

Parameter: γ_2

with ± 1.96 standard deviations




Figure 10: Parameter γ_2 bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: w_1

with ± 1.96 standard deviations




Figure 11: Parameter w_1 bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: w_2

with ± 1.96 standard deviations




Figure 12: Parameter w_2 bias with ± 1.96 standard deviations. cs means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $\log(\sigma_1^2)$

with ± 1.96 standard deviations




Figure 13: Parameter $\log(\sigma_1^2)$ bias with ± 1.96 standard deviations. cs means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $\log(\sigma_2^2)$

with ± 1.96 standard deviations




Figure 14: Parameter $\log(\sigma_2^2)$ bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $\log(\sigma_3^2)$

with ± 1.96 standard deviations




Figure 15: Parameter $\log(\sigma_3^2)$ bias with ± 1.96 standard deviations. *cs* means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $\log(\sigma_4^2)$

with ± 1.96 standard deviations




Figure 16: Parameter $\log(\sigma_4^2)$ bias with ± 1.96 standard deviations. cs means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $z(\rho_{12})$

with ± 1.96 standard deviations




Figure 17: Parameter $z(\rho_{12})$ bias with ± 1.96 standard deviations. cs means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Parameter: $z(\rho_{34})$
with ± 1.96 standard deviations




Figure 18: Parameter $z(\rho_{34})$ bias with ± 1.96 standard deviations. **cs** means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

Complete model's cross-correlations

with ± 1.96 standard deviations




Figure 19: Parameter $\{z(\rho_{13}), z(\rho_{24}), z(\rho_{14}), z(\rho_{23})\}$ bias with ± 1.96 standard deviations. **cs** means the cluster size, and 5-30-60k the sample size. Bar width is given by the number of converged fittings.

References

- Andersen, P. K., Geskus, R. B., de Witte, T. and Putter, H. (2012). Competing risks in epidemiology: possibilities and pitfalls, International Journal of Epidemiology **31**(1): 861–870.
- Bonat, W. H. (2018). Multiple Response Variables Regression Models in R: The `mcglm` Package, Journal of Statistical Software **84**(4).
- Bonat, W. H. and Jørgensen, B. (2016). Multivariate covariance generalized linear models, Journal of the Royal Statistical Society, Series C (Applied Statistics) **65**(5): 649–675.
- Bonat, W. H. and Ribeiro Jr, P. J. (2016). Practical likelihood analysis for spatial generalized linear mixed models, Environmetrics **27**(1): 83–89.
- Breslow, N. E. and Clayton, D. G. (1993). Approximate inference in generalized linear mixed models, Journal of the American Statistical Association **88**(421): 9–25.
- Cederkvist, L., Holst, K. K., Andersen, K. K. and Scheike, T. H. (2019). Modeling the cumulative incidence function of multivariate competing risks data allowing for within-cluster dependence of risk and timing, Biostatistics **20**(2): 199–217.
- Cheng, Y. and Fine, J. P. (2012). Cumulative incidence association models for bivariate competing risks data, Journal of the Royal Statistical Society, Series B (Methodological) **74**(2): 183–202.
- Cheng, Y., Fine, J. P. and Kosorok, M. R. J. (2007). Nonparametric Association Analysis of Bivariate Competing-Risks Data, Journal of the American Statistical Association **102**(480): 1407–1415.
- Cheng, Y., Fine, J. P. and Kosorok, M. R. J. (2009). Nonparametric Association Analysis of Exchangeable Clustered Competing Risks Data, Biometrics **65**(1): 385–393.
- Clayton, D. G. (1978). A model for association in bivariate life tables and its application in epidemiological studies of familial tendency in chronic disease incidence, Biometrika **65**(1): 141–151.
- Cox, D. R. and Reid, N. (2004). A note on pseudolikelihood constructed from marginal densities, Biometrika **91**(3): 729–737.
- Dennis, J. E., Gay, D. M. and Welsch, R. E. (1981). An Adaptive Nonlinear Least-Squares Algorithm, ACM Transactions on Mathematical Software **7**(3): 348–368.
- Diaconis, P. (2009). The Markov chain Monte Carlo revolution, Bulletin (New Series) of the American Mathematical Society **46**(2): 179–205.

- 1 Embrechts, P. (2009). Copulas: A Personal View, The Journal of Risk and Insurance
2 **76**(3): 639–650.
- 3
- 4 Fine, J. P. (1999). Analysing competing risks data with transformation models, Journal
5 of the Royal Statistical Society, Series B (Methodological) **61**(4): 817–830.
- 6
- 7
- 8 Fine, J. P. and Gray, R. J. (1999). A proportional hazards models for the subdistribution
9 of a competing risk, Journal of the American Statistical Association **94**(446): 496–509.
- 10
- 11
- 12 Gay, D. M. (1990). Usage summary for selected optimization routines, Technical report,
13 Computing Science Technical Report 153, AT&T Bell Laboratories. Murray Hill, NJ.
- 14
- 15
- 16 Gelfand, A. E. and Smith, A. F. M. (1990). Sampling-Based Approaches to Calculating
17 Marginal Densities, Journal of the American Statistical Association **85**(410): 398–409.
- 18
- 19
- 20 Gerds, T. A., Scheike, T. H. and Andersen, P. K. (2012). Absolute risk regression for
21 competing risks: interpretation, link functions and prediction, Statistics in Medicine
22 **31**(29): 3921–3930.
- 23
- 24
- 25
- 26 He, Y., Kim, S., Mao, L. and Ahn, K. W. (2022). Marginal semiparametric transformation
27 models for clustered multivariate competing risks data, Statistics in Medicine **41**: 5349–
28 5364.
- 29
- 30
- 31 Hoffman, M. D. and Gelman, A. (2014). The No-U-Turn Sampler: Adaptively Setting
32 Path Lengths in Hamiltonian Monte Carlo, Journal of Machine Learning Research
33 **15**(47): 1593–1623.
- 34
- 35
- 36
- 37 Hougaard, P. (2000). Analysis of Multivariate Survival Data, Springer-Verlag, New York.
- 38
- 39
- 40 Kalbfleisch, J. D. and Prentice, R. L. (2002). The Statistical Analysis of Failure Time
41 Data, second Edition edn, John Wiley & Sons, Inc., Hoboken, New Jersey.
- 42
- 43
- 44 Klein, J. P. (1992). Semiparametric estimation of random effects using cox model based
45 on the em algorithm, Biometrics **48**(1): 795–806.
- 46
- 47
- 48 Kristensen, K., Nielsen, A., Berg, C. W., Skaug, H. J. and Bell, B. M. (2016). TMB:
49 Automatic Differentiation and Laplace Approximation, Journal of Statistical Software
50 **70**(5): 1–21.
- 51
- 52
- 53 Krupskii, P. and Joe, H. (2013). Factor copula models for multivariate data, Journal of
54 Multivariate Analysis **120**(1): 85–101.
- 55
- 56
- 57 Kuk, A. Y. C. (1992). A semiparametric mixture model for the analysis of competing
58 risks data, Australian Journal of Statistics **34**(2): 169–180.
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1 Laird, N. M. and Ware, J. H. (1982). Random-effects models for longitudinal data,
2 Biometrics **38**(4): 963–974.
- 3
4 Larson, M. G. and Dinse, G. E. (1985). A Mixture Model for the Regression Analysis
5 of Competing Risks Data, Journal of the Royal Statistical Society, Series C (Applied
6 Statistics) **34**(3): 201–211.
- 7
8
9 Liang, K. Y., Self, S., Bandeen-Roche, K. J. and Zeger, S. L. (1995). Some recent develop-
10 ments for regression analysis of multivariate failure time data, Lifetime Data Analysis
11 **1**(1): 403–415.
- 12
13
14 Lindsay, B. G. (1988). Composite likelihood methods, Comtemporary Mathematics
15 **80**(1): 221–239.
- 16
17
18 Masarotto, G. and Varin, C. (2012). Gaussian copula marginal regression, Electronic
19 Journal of Statistics **6**(1): 1517–1549.
- 20
21
22 McCullagh, P. and Nelder, J. A. (1989). Generalized linear models, second edition edn,
23 Chapman & Hall, London.
- 24
25
26 McCulloch, C. E. and Searle, S. R. (2001). Generalized, Linear, and Mixed Models, John
27 Wiley & Sons, Inc., New York.
- 28
29
30 Molenberghs, G. and Verbeke, G. (2005). Models for Discrete Longitudinal Data, Springer,
31 New York.
- 32
33
34 Monnahan, C. and Kristensen, K. (2018). No-U-turn sampling for fast Bayesian inference
35 in ADMB and TMB: Introducing the `adnuts` and `tmbstan` R packages, PloS ONE
36 **13**(5).
- 37
38
39 Naskar, M., Das, K. and Ibrahim, J. G. (2005). A Semiparametric Mixture Model for
40 Analyzing Clustered Competing Risks Data, Biometrics **61**(3): 729–737.
- 41
42
43 Nelder, J. A. and Wedderburn, R. W. M. (1972). Generalized linear models, Journal of
44 the Royal Statistical Society, Series A **135**(3): 370–384.
- 45
46
47 Nielsen, G. G., Gill, R. D., Andersen, P. K. and Sørensen, T. I. A. (1992). A Counting
48 Process Approach to Maximum Likelihood Estimation in Frailty Models, Scandinavian
49 Journal of Statistics **19**(1): 25–43.
- 50
51
52 Nocedal, J. and Wright, S. J. (2006). Numerical Optimization, Springer Series in Opera-
53 tions Research and Financial Engineering, second Edition edn, Springer, New York.
- 54
55
56 Petersen, J. H. (1998). An Additive Frailty Model for Correlated Life Times, Biometrics
57 **54**(1): 646–661.
- 58
59
60
61
62
63
64
65

- Peyré, G. (2020). Course notes on optimization for machine learning, May 10, <https://mathematical-tours.github.io/book-sources/optim-ml/OptimML.pdf>. CNRS & DMA, École Normale Supérieure.
- Prentice, R. L., Kalbfleisch, J. D., Peterson Jr, A. V., Flournoy, N., Farewell, V. T. and Breslow, N. E. (1978). The analysis of failure times in the presence of competing risks, *Biometrics* **1**(1): 541–554.
- R Core Team (2021). *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing. Vienna, Austria. <https://www.R-project.org/>.
- Scheike, T. and Sun, Y. (2012). On cross-odds ratio for multivariate competing risks data, *Biostatistics* **13**(4): 680–694.
- Scheike, T., Zhang, Y. S. M. and Jensen, T. K. (2010). A semiparametric random effects model for multivariate competing risks, *Biometrika* **97**(1): 133–145.
- Shi, H., Cheng, Y. and Jeong, J. H. (2013). Constrained parametric model for simultaneous inference of two cumulative incidence functions, *Biometrical Journal* **55**(1): 82–96.
- Shih, J. H. and Albert, P. S. (2009). Modeling Familial Association of Ages at Onset of Disease in the Presence of Competing Risk, *Biometrics* **66**(4): 1012–1023.
- Shun, Z. and McCullagh, P. (1995). Laplace approximation of high dimensional integrals, *Journal of the Royal Statistical Society, Series B (Methodological)* **57**(4): 749–760.
- Stan Development Team (2019). *Stan Modeling Language Users Guide and Reference Manual*, Version 2.26. <https://mc-stan.org>.
- Stan Development Team (2020). RStan: the R interface to Stan, <https://mc-stan.org/>. R package version 2.21.2.
- Therneau, T. M. and Grambsch, P. M. (2000). *Modeling Survival Data: Extending the Cox Model*, Springer-Verlag, New York.
- Tierney, L. and Kadane, J. (1986). Accurate approximations for posterior moments and marginal densities, *Journal of the American Statistical Association* **81**(393): 82–86.
- Varin, C., Reid, N. and Firth, D. (2011). An overview of composite likelihood methods, *Statistica Sinica* **21**(1): 5–42.
- Vaupel, J. W., Manton, K. G. and Stallard, E. (1979). The impact of heterogeneity in Individual Frailty on the Dynamics of Mortality, *Demography* **16**(1): 439–454.
- Wood, S. N. (2015). *Core Statistics*, Institute of Mathematical Statistics, Textbooks, IMS.